




# HIV IN MARYLAND, 2016

OCTOBER 2017

## New HIV Diagnoses


- In 2016, there were **1,118** adult/adolescent HIV cases (age 13+) diagnosed in Maryland. Baltimore City and Prince George's County had the highest rates of new HIV diagnoses.
- Proportion of HIV diagnoses in Maryland has **decreased** in men from **84.7% in 1985** to **72.2% in 2016**, and has **increased** in women from **15.3% in 1985** to **27.8% in 2016**.

## Trends in HIV Diagnoses by Gender



- 73.7%** of the **1,118** reported HIV diagnoses (age 13+) in Maryland in 2016 were **NH-Black**, **13.6%** were **NH-White**, **8.2%** were Hispanic, and **4.5%** were **NH-Other** races.
- Of the **1,118** adult/adolescent HIV diagnoses (age 13+) in 2016, **35.2%** were among adults ages **20-29**, **23.9%** were among those ages **30-39**, and **17.1%** were among those ages **40-49**.

## Maryland Adult/Adolescent HIV Diagnoses, Rates by Jurisdiction, 2016



## Characteristics of Marylanders Living with HIV


### Exposure Category\*

- 38.8% MSM**
- 37.8% HET**
- 19.6% IDU**
- 3.6% MSM/IDU**
- 0.2% Other**

### Race/Ethnicity


- 74.2% NH-Black**
- 13.1% NH-White**
- 6.3% Hispanic**
- 5.8% NH-Other**

- Case rates (per 100,000) were **twice** as high in **males (836.2)** compared to **females (387.5)**. In other words, **1 in 119 males** and **1 in 258 females** in Maryland were living with HIV.



- Total living HIV case rates in Maryland were **3 to 10 times higher** among NH-Blacks (**1,520.1**) compared to Hispanics (**436.9**) and NH-Whites (**148.0**).

## Living HIV Case Rates by Race/Ethnicity



**At the end of 2016, there were 30,430 living adults/adolescents diagnosed with HIV in Maryland.**

**HET:** Heterosexual Exposure | **IDU:** Injection Drug User | **MSM:** Men who have Sex with Men | **NH:** Non-Hispanic

Center for HIV Surveillance, Epidemiology and Evaluation  
Prevention and Health Promotion Administration  
Maryland Department of Health

\* Multiple imputation was used to estimate and adjust for missing transmission category.  
Data reported through 06/30/2017 from the Enhanced HIV/AIDS Reporting System.